



New records of *Ganoderma multiplicatum* (Mont.) Pat. (Polyporales, Basidiomycota) from Colombia and its geographic distribution in South America

Ana Cristina Bolaños^{1, 2*}, Vera Lucia Ramos Bononi², Adriana de Mello Gugliotta² and Jaime Eduardo Muñoz³

1 Universidad del Valle, Departamento de Biología. Calle 13 No. 100–00, Cali, Colombia

2 Instituto de Botânica de São Paulo. Núcleo de Pesquisa em Micologia. Av. Miguel Stéfano, 3687, SP, Brazil

3 Universidad Nacional de Colombia, Departamento de Ciencias Biológicas. Carrera 32 No. 12–00, Palmira, Colombia

* Corresponding author. E-mail: ana.bolanos@correounivalle.edu.co

Abstract: *Ganoderma* is a cosmopolitan genus of fungi with species distributed in temperate and tropical regions. Species of *Ganoderma* in living Leguminosae were observed in Park de la Salud in Pance, Cali, Colombia and we record *G. multiplicatum* for the first time from Colombia. A distribution map of this genus in the Neotropical region is presented.

Key words: Ganodermataceae; Leguminosae; taxonomy; Neotropics

The genus *Ganoderma* P. Karst was created by Karsten (1881) to recognize a lacquered and stipitate fungus, *Polyporus lucidus* (Curtis) Fr., the type of the genus (Moncalvo and Ryvarden 1997). *Ganoderma* is well circumscribed by its truncated, ellipsoid basidiospore with apical germinal pore and, double walls structure; the external wall (exosporium) is thin and hyaline, and the internal wall (endosporium) is dark brown and separated with internal crests (Ryvarden 2004). Due to the high phenotypic plasticity found in the basidiomata of this genus and varying concepts of the species, identifying some *Ganoderma* species is confusing and correct geographical distributions are still pending (Moncalvo and Ryvarden 1997). The genus is considered to be cosmopolitan with species distributed in temperate and tropical regions worldwide (Cao and Yuan 2013). In Colombia, *Ganoderma* has not been studied in depth, but morphology-based studies and previous work on the diversity of macrofungi in Colombia included thirteen species of *Ganoderma* (Vasco-Palacios and Franco-Molano 2013). A list of species, locality, voucher and references are presented in Table 1.

Ganoderma multiplicatum (Mont.) Pat. was originally collected in French Guyana on a tree trunk (Moncalvo and Ryvarden 1997). This species has been considered most similar to *G. chalceum* (Corner 1983). However, Ryvarden (2000) considered it a synonym of *G. subamboinense* Henn., but a molecular phylogenetic analysis (ITS region) by Lima et al. (2014) showed that *G. chalceum* and *G. subamboinense* are in different clades, suggesting they are not synonymous. During a search for species of *Ganoderma* in living Leguminosae and in rotten logs in urban and rural areas, we collected a basidiome of *G. multiplicatum* at Park de la salud (03°19'42" N, 076°38'19" W), Cali, Colombia (Table 1). The collection was made according to the methodology of Fidalgo and Bononi (1984). To analyze morphological characters, thin sections of basidiome were prepared and treated with 4% KOH, 1% Phloxine, and Melzer reagent (Ryvarden 2004). Vouchers were deposited at the herbarium of the Botanical Institute of São Paulo (SP) and the herbarium at the Valley University in Colombia (CUVC). DNA analysis (isolation, amplification and sequencing) for ITS region was performed according to protocols used by Raeder and Broda (1985), Moncalvo et al. (1995), Gottlieb et al. (2000). Primers set BMB-CR (5'-GTACACACCGCCCGTCG-3') and LRI (5'-GGTTGGTTTCTTTTCCT-3') were used under conditions described by Vilgalys and Hester (1990). Sequences were aligned using BioEdit 7.2.0 (Hall 1999). Best model of nucleotide substitution (HKY+G) was selected by the Akaike criterion in JModelTest version 2.1.3 (Darriba et al. 2012). Maximum parsimony and likelihood was performed using PAUP version 4.0b10 (Swofford 2002). Bayesian analysis was performed using MrBayes 3.04b (Ronquist and Huelsenbeck 2003).

Table 1. Colombian species of *Ganoderma*, locality, voucher and references.

Species	Department/ locality Voucher	Reference
<i>G. adpersum</i> (Schulzer) Donk	Cundinamarca. Loc. n. det. AR0019, AR 0049	Ruiz and Varela (2006)
<i>G. amazonense</i> Weir	Cundinamarca, Zipacón. Ruiz, A. HPUJ 0031 Chocó. Loc. n. det., Torres 131 Chocó, Las animas. CO-5788	Ruiz and Varela (2006) Guzmán et al. (2004) Setliff and Ryvarden (1983)
<i>G. applanatum</i> (Pers.) Pat.	Caldas, Parque los Nevados. Loc. and Vouch. n. det. Dept., Loc. and Vouch. n. det. Caldas, Parque los Nevados, COLTB827 Caquetá, Solano. Vasco-P, A. 529 Caquetá, Peña Rioja. López-Q. 265 Meta, San Juan de Arama, CO-2425	Boekhout and Pulido (1989) Dennis (1970) Pulido and Boekhout (1989) Vasco-Palacios et al. (2005) Vasco-Palacios et al. (2005) Setliff and Ryvarden (1983)
<i>G. applanatum</i> var. <i>tornatum</i> (Pers.) C.J. Humphrey	Dept., Loc. and Vouch. n. det. Caquetá, Florencia-El Doncello road. CO-3042	Dennis (1970) Setliff and Ryvarden (1983)
<i>G. australe</i> (Fr.) Pat.	Amazonas, La Chorrera. Vasco-P, A. 218 Cundinamarca, Chia city. Pinzón-O, C. HUPN 055, 065, 081, 090, 105, 110, 130, 155, 160, 179 and 190 Chocó. Loc. n. det., Torres 95 Chocó. Loc. and Vouch. n. det.	Vasco-Palacios et al. (2005) Pinzón-Osorio and Pinzón-Osorio (2016) Guzmán et al. (2004) Torres and Hurtado(2003)
<i>G. brownii</i> (Murrill) Gib.	Cundinamarca, Cerro de las Mercedes Guzmán 4639	Guzmán and Varela (1978)
<i>G. coffeatum</i> (Berk.) J.S. Furtado	Boyacá, Chocontá-Agua Clara road. CO-4762 Chocó, Quibdó-Itzmina road. CO-5680, CO-5683 Chocó, Loc. n. det., Guzmán 35321	Setliff and Ryvarden (1983) Setliff and Ryvarden (1983) Guzmán et al. (2004)
<i>G. concinnum</i> Ryvarden	Chocó, Riosucio nat. park. Ryvarden 16840 (O, Holotype, NY, Isotype)	Ryvarden (2004), Ryvarden (2000)
<i>G. chalceum</i> (Cooke) Steyaert	Vaupés, Tio Barbas, Vaupés River. Holliday PH51(Holotype, K)	Ryvarden (2004), Ryvarden (2000)
<i>G. fornicatum</i> (Fr.) Pat.	Dept., Loc. and Vouch. n. det.	Dennis (1970)
<i>G. lucidum</i> (Curtis) P. Karst.	Caquetá, Solano. Alvarez 32, López-Q. 51 Caquetá. Loc. n. det., Vouch. n. det. Cundinamarca, Colorado. Guzmán 4539 Chocó. Loc. n. det., Guzmán 35407 Chocó. Loc. n. det., Vouch. n. det. Valle del Cauca, Cali city. Guzmán 4473	Vasco-Palacios et al. (2005), Franco-Molano et al. (2005) Guzmán and Varela (1978) Guzmán et al. (2004) Torres and Hurtado(2003) Guzmán and Varela (1978)
<i>G. nitidum</i> Murrill	Caquetá. Loc. n. det., Vouch. n. det. Caquetá, Solano. Vasco-P, A. HUA139	Franco-Molano et al. (2005) Vasco-Palacios et al. (2005)
<i>G. tornatum</i> (Pers.) Bres.	Cundinamarca, El bosque de Tibabita. CO-14	Setliff and Ryvarden (1983)

Loc. n. det. = local no determinate, Vouch. n. det. = Voucher no determinate, Dept., Loc. and Vouch. n. det. = department, local and voucher no determinate.

***Ganoderma multiplicatum* (Mont.) Pat., Bull. Soc.**

Mycol. Fr. 5: 74, 1889.

≡ *Polyporus multiplicatus* Mont., Ann. Sci. Nat. Bot. Ser. 4, 1: 128, 1854.

Figures 1–9.

This species presents basidiomata 14 × 10 × 3 cm, dimidiate to sessile perennial, margin whitish, lobulate, shiny laccate surface, pale light brown to reddish brown, pores angular to round, 4–6 per mm, pore surface creamy to ochraceous; cuticle formed by clavate hyphal ends, strongly amyloid in the apical part, with a few protuberances and basidiospores 8.4 × 5.8 μm.


Specimen examined: Colombia: Cali, Dept. Valle, Park de la Salud, Pance, 05.I.2013. A.C. Bolaños and G. Aguilar (CUVC60077).

Additional specimens examined: Brazil. São Paulo, SP417797, SP417778, SP417780; French Guyana K123639 lectotype.

In the Neotropical region, *Ganoderma multiplicatum* was cited in Venezuela (Ryvarden 2004) and in Brazil it has been mentioned from Alagoas (Torres-Torres et al. 2012), Amazonas, Para and Rondônia (Gomes-Silva and

Gibertoni 2009; Torres-Torres et al. 2012), Mato Grosso do Sul (Quevedo et al. 2012), Paraíba (Lima et al. 2014), Rio de Janeiro (Torres-Torres et al. 2012), Roraima (Gomes-Silva et al. 2011), São Paulo (Gugliotta et al. 2011; Torres-Torres et al. 2012), Santa Catarina (Loguercio-Leite et al. 2005; Drechsler-Santos et al. 2008; Baltazar and Gibertoni 2009), and Sergipe (Torres-Torres et al. 2012). In Argentina, this species has been reported from Tucumán, Corrientes and Chaco (Gottlieb and Wright 1999). A list of localities, substrates and vouchers can be found in Table 2.

This is the first record of the species from Colombia and contributes to the list of *Ganoderma* species from Colombia and the Neotropics. The type material of *G. multiplicatum*, from French Guyana, is in Kew herbarium (K), and it consists of a basidiome fragment. Nevertheless, it allowed for the identification of our Colombian specimen. Although originally described from South America, *G. multiplicatum* was subsequently found in Africa and Asia (Steyaert 1980; Zhao 1989), including India (Bhosle et al. 2010) and Taiwan (Wang



Figures 1–6. *Ganoderma multiplicatum*. **1.** Basidiome. **2.** Context. **3.** Cuticle cell. **4.** Pores. **5–6.** Basidiospores. Scale bar 1 = 2 cm, 2 = 1 mm, (3–5) = 10 μm, 4 = 100 μm, 6 = 2 μm.

Table 2. Geographic distribution, ecosystem, substrate and references where *Ganoderma multiplicatum* has been found in America.

Country Department/state, locality Voucher	Ecosystem, substrate	Reference
Argentina		
Corrientes, Corrientes city, BACF 34461	On <i>Schinopsis</i> sp. stump	Gottlieb and Wright (1999)
Corrientes, Mbucuruyá, Sta Teresa Station, BAFC 24414	Tipuana tipu	Gottlieb and Wright (1999)
Corrientes, Cosme Dept. BAFC 24416	Buried wood	Gottlieb and Wright (1999)
Chaco, Charata	N.d.*	Gottlieb and Wright (1999)
Tucuman, Tucuman city, BAFC 34390	On stump	Gottlieb and Wright (1999)
Tucuman, La Cocha Dept., Monte Grande, Lil.8846	N.d.*	Gottlieb and Wright (1999)
Brazil		
Alagoas, Maceió Ibama Reserve. Maziero, R. s.n. SP250605	N.d.*	Torres-Torres et al. (2012)
Amazonas, Santa Laura, Madeira River. U.S. dept. Agric. Path. Myc. Coll. 66859, s.d. RLS.55.K.25	N.d.*	Steyaert (1980)
Amazonas, Manaus. M.A. Sousa 151.INPA74707	N.d.*	Gomes-Silva and Gilbertoni (2009), Torres-Torres et al. (2012)
Amazonas, Manaus. M.A. Sousa 249.INPA74533	N.d.*	Gomez-Silva et al. (2011)
Amazonas, Barcelos, Serra de Aracá, E.S.S. et al. 294. INPA 153737	N.d.*	Gomez-Silva et al. (2011)
Mato Grosso do Sul, Campo grande, mata do prosa. J.R. Quevedo & A.K.M. Oliveira 11SP	Living trees	Quevedo et al. (2012)
Para, Melgaço, scientific station Ferreira penna. T.B. Gilbertoni URM 79280, URM79281, URM79282, URM79283	N.d.*	Gomes-Silva et al. (2011)
Rio de Janeiro, mata da Tijuca. J.S. Furtado s.n. SP91404	N.d.*	Torres-Torres et al. (2012)
Paraíba, Mataracá, Mineradora Millenium. URM83346	N.d.*	Lima et al. (2014)
Rondônia, Jaru, right margin of Jaru river. Capelari & Maziero, 693. SP211484	N.d.*	Gomes-Silva et al. (2011); Torres-Torres et al. (2012)
Rondônia, Loc. n. det., R.D. Goos et al. 1673. INPA125007	N.d.*	Gomes-Silva et al. (2011)
Rondônia, Porto Velho, municipal natural park de Porto Velho. A.C. Gómez-Silva 646. URM81081.	N.d.*	Gomes-Silva et al. (2011), Lima et al. (2014)
Roraima, Boa vista, M.A. Jesús 882, INPA186187	N.d.*	Gómez-Silva et al. (2011)

Continued

Table 1. Continued.

Country Department/state, locality Voucher	Ecosystem, substrate	Reference
Santa Catarina, Loc. n. det.	N.d.*	Drechsler-Santos et al. (2008), Baltazar and Gibertoni (2009),
Santa Catarina, Florianópolis, Leite & Folle. (FLOR10841). As G. subamboinense.	N.d.*	Loguercio-Leite et al. (2005), Drechsler-Santos et al. (2008), Baltazar and Gilbertoni (2009)
Santa Catarina, Florianópolis. Folle & Willerding. (FLOR10859). As G. subamboinense	N.d.*	Loguercio-Leite et al. (2005)
São Paulo, Parque Estadual Fontes do Ipiranga. Campacci 12(SP)	N.d.*	Gugliotta et al. (2011), Torres-Torres et al. (2012)
São Paulo, Parque do estado, Botanical Institute. B. Skovortzov s.n. SP10746	On Wood	Torres-Torres et al. (2012)
Sergipe, Ecological station Santa Isabel. R.H. Marino s.n. SP375895	N.d.*	Torres-Torres et al. (2012)
Colombia		
Cali, Parque de la Salud. A.C. Bolaños & G. Aguilar 294. CUV60077	Caesalpinia pelthophoroides	This work
French Guiana		
Isla Principe, Loc. n. det., Crypt. Guyan. 357	On trunk	Steyaert (1980); Moncalvo and Ryvarden (1997)
Venezuela		
Loc. n. det.	N.d.*	Ryvarden (2004)

N.d.* = no data, Loc. n. det. = local no determinate.




Figure 7–9. *Ganoderma multiplicatum*. 7. Basidiospores. 8. Cuticle cells. 9. Skeletal hyphae. Scale bar = 5 μm.




Figure 10. Geographical distribution map of *Ganoderma multiplicatum* from Colombia and South America. (Source map: Robledo et al. 2016).

and Wu 2007). The distribution of *G. multiplicatum* in the Neotropics is shown in Figure 10.

In our phylogenetic analysis, the Colombian sample as was found to be *G. multiplicatum*, with high statistical support (Bootstrap: MP: 80%, ML: 82%, BPP: 0.9), using the ITS region and including 26 additional sequences from GenBank. According to Lima et al. (2014), and in this analysis, *G. chalceum* and *G. subamboinense* were recovered in different clades (Figure 11).

ACKNOWLEDGEMENTS

We thank SP and CUVIC herbaria for facilitating the equipment used in this study. We also thank the Support Foundation for Scientific Research of the State of São Paulo (FAPESP) and the Universidad del Valle in Colombia for financial support, Alex Mora for his help with the molecular analyses in PAUP program, Viviana Motato for the illustration, and Dr. Gerardo Robledo for the map.




Figure 11. ITS tree and reconstruction of the phylogenetic ITS of the sequence of *G. multiplicatum*. Statistical support nodes include the maximum parsimony bootstrap, maximum likelihood and Bayesian respectively.

LITERATURE CITED

- Baltazar, J.M. and T.B. Gibertoni. 2009. A checklist of aphyllophoroid fungi (Basidiomycota) recorded from the Brazilian Atlantic rain forest. *Mycotaxon* 109: 439–442 <http://www.mycotaxon.com/resources/checklists/baltazar-v109-checklist.pdf>
- Boekhout, T. and M.M. Pulido. 1989. The occurrence of macrofungi and their habitats in vegetations along The Parque Los Nevados transect; pp. 507–515, in: T. Van der Hammen, S. Diaz-Piedrahita and J. Alvarez (eds.). *La cordillera central colombiana transecto Parque Los Nevados (segunda parte)*. Berlin: J. Cramer; *Estudios de Ecosistemas Tropoandinos*, 3.
- Bhosle, S., K. Ranadive, G. Bapat, S. Garad, G. Deshpande and J. Vaidya. 2010. Taxonomy and diversity of *Ganoderma* from the western parts of Maharashtra (India). *Mycosphere* 1: 249–262 http://www.mycosphere.org/pdf/MC1_3_No8.pdf
- Corner, E.J.H. 1983. *Ad Polyporaceas I. Amauroderma and Ganoderma*. *Beihefte zur Nova Hedwigia* 75: 1–182.
- Dariva, D., G.L. Taboada, R. Doallo and D. Posada. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772–772.
- Dennis, R.W. 1970. *Fungus flora of Venezuela and adjacent countries*. Kew Bulletin, additional series III. London: J. Cramer. Vaduz. 531 pp.
- Drechsler-Santos, E.R., C. Groposo and C. Loguercio-Leite. 2008. Additions to the knowledge of lignocellulolytic basidiomycetes in forests from Santa Catarina, Southern Brazil. *Mycotaxon* 103: 197–200. <http://www.mycotaxon.com/resources/checklists/drechsler-v103-checklist.pdf>
- Fidalgo, O. and V.R.L. Bononi. 1984. *Técnicas de coleta, preservação e herborização de material botânico*. São Paulo: Instituto de Botânica. 62 pp.
- Gomes-Silva, A.C. and T.B. Gibertoni. 2009. Checklist of the aphyllophoraceous fungi (Agaricomycetes) of the Brazilian Amazonia. *Mycotaxon* 108: 319–322. <http://www.mycotaxon.com/resources/checklists/gomesSilva-v108-checklist.pdf>
- Gomes-Silva, A.C., L. Ryvarden and T.B. Gibertoni. 2011. New records of Ganodermataceae (Basidiomycota) from Brazil. *Nova Hedwigia* 92(1–2): 83–94. doi: 10.1127/0029-5035/2011/0092-0083
- Gottlieb, A.M. and J.E. Wright. 1999. Taxonomy of *Ganoderma* from southern South America: subgenus *Ganoderma*. *Mycological Research* 103: 1289–1289.
- Gottlieb, A.M., E. Ferrer and J.E. Wright. 2000. rDNA analyses as an aid to the taxonomy of species of *Ganoderma*. *Mycological Research* 104: 1033–1045.
- Gugliotta, A.M., G.D. Poscolere and T.V. Silva. 2011. Criptógamos do Parque Estadual das Fontes do Ipiranga, São Paulo, SP, Brasil. *Fungos*, 10: Ganodermataceae. *Hoehnea* 38(4): 687–695 <http://www.scielo.br/pdf/hoehnea/v38n4/09.pdf>
- Hall, T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98. <http://brownlab.mbio.ncsu.edu/JWB/papers/1999Hall1.pdf>
- Karsten, P.A. 1881. *Enumeratio boletinearum et polyporearum fennicarum, systemate novo dispositarum*. *Revue Mycologique Toulouse* 3: 16–19.
- Lima, N.J., G. Baptista and E. Malosso. 2014. Delimitation of some Neotropical laccate *Ganoderma* (Ganodermataceae): molecular phylogeny and morphology. *Revista de Biología Tropical* 62(3): 1197–1208. <http://revistas.ucr.ac.cr/index.php/rbt/article/view/12380/14744>
- Loguercio-Leite C., C. Groposo and M.A. Halmenschlager. 2005. Species of *Ganoderma* Karsten in a subtropical area (Santa Catarina state, Southern Brazil). *Iheringia, série Botânica* 60: 135–139.
- Moncalvo, J.M. and L. Ryvarden. 1997. A nomenclatural study of the Ganodermataceae Donk. *Synopsis Fungorum* 11. Oslo: Fungiflora. 114 pp.

- Moncalvo, J.M., H.H. Wang and R.S. Hseu. 1995. Phylogenetic relationships in *Ganoderma* inferred from the internal transcribed spacers and 25S ribosomal DNA sequences. *Mycologia* 87(2): 223–238. <http://www.jstor.org/stable/3760908>
- Pinzón-Osorio, C.A. and J. Pinzón-Osorio. 2016. Ampliación de la distribución geográfica de *Ganoderma australe* (Patouillard, 1889) (Aphyllophorales-Basidiomycota), para el departamento de Cundinamarca, Colombia. *Revista Biodiversidad Neotropical* 6: 7–11. doi: 10.18636/bioneotropical.v6i1.180
- Pulido, M.M. and T. Boekhout. 1989. Distribution of macrofungi along the Parque Los Nevados transect; pp. 485–505, in: T. Van der Hammen, S. Díaz-Piedrahita and V.J. Alvarez (eds.). La cordillera central colombiana transecto Parque Los Nevados. (segunda parte). *Studies on Tropical Andean Ecosystems* 3.
- Quevedo, J., V.L. Bononi, A.K.M. Oliveira and A.M. Gugliotta. 2012. Agaricomycetes (Basidiomycota) em um fragmento florestal urbano na cidade de Campo Grande, Mato Grosso do Sul, Brasil. *Revista Brasileira de Biociências* 10: 430–438. <http://www.ufrgs.br/seerbio/ojs/index.php/rbb/article/view/1998/1166>
- Raeder, U. and P. Broda. 1985. Rapid preparation of DNA from filamentous fungi. *Letters in Applied Microbiology* 1(1): 17–20. doi: 10.1111/j.1472-765X.1985.tb01479.x
- Robledo, G.L., E.M. Giorgio, C.R.P. Franco, O. Popoff and C. Decock. 2014. *Gyrodontium sacchari* (Spreng. Fr.) Hjortstam (Boletales, Basidiomycota) in America. New records and its geographic distribution. *Check List* 10(6): 1514–1519. doi: 10.15560/10.6.1514
- Ronquist, F. and J.P. Huelsenbeck, 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Ruiz, A. and A. Varela. 2006. Nuevos registros de Aphyllophorales (Basidiomycota) en bosque montano húmedo y de niebla de Colombia. *Caldasia* 28(2): 259–266. <http://www.scielo.org.co/pdf/cal/v28n2/v28n2a9.pdf>
- Ryvarden, L. 2000. Studies in Neotropical polypores 2: a preliminary key to Neotropical species of *Ganoderma* with a laccate pileus. *Mycologia* 92: 180–191.
- Ryvarden, L. 2004. Neotropical polypores: Part 1. Introduction, Ganodermataceae & Hymenochataceae. Oslo: Fungiflora. 227 pp.
- Setliff, E.C. and L. Ryvarden. 1983. Los hongos de Colombia 7. Some Aphyllophorales wood inhabiting fungi. *Mycotaxon* 18(2): 509–525.
- Swofford, D.L. 2002. PAUP*. Phylogenetic Analysis Using Parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland.
- Steyaert, R.L. 1980. Study of some *Ganoderma* species. *Bulletin du Jardin Botanique National de Belgique* 50: 135–186.
- Torres-Torres, M.G., L. Guzmán-Dávalos and A.M. Gugliotta. 2012. *Ganoderma* in Brazil. *Mycotaxon* 121: 93–132.
- Torres, G. and A.L. Hurtado. 2003. Micobiota nativa con potencialidades medicinal en el Municipio de Quibdó. *Revista Institucional, Universidad Tecnológica del Chocó: Investigación, Biodiversidad y Desarrollo* 18: 3–9.
- Vasco-Palacios, A.M. and A.E. Franco-Molano. 2013. Diversity of Colombian macrofungi (Ascomycota-Basidiomycota). *Mycotaxon* 121: 499. <http://www.mycotaxon.com/resources/checklists/VascoPalacios-v121-checklist.pdf>
- Vasco-Palacios, A.M., A.E. Franco-Molano, C.A. López-Quintero and T. Boekhout. 2005. Macromicetes (ascomycota, basidiomycota) de la región del medio Caquetá, departamentos de Caquetá y Amazonas (Colombia). *Biota Colombiana* 6(1): 127–140. <http://www.redalyc.org/pdf/491/49106101.pdf>
- Wong, D.M. and Sheng-Hua, W. 2007. Two species of *Ganoderma* new to Taiwan. *Mycotaxon* 102: 373–378.
- Vilgalys, R. and M. Hester. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172(8): 4238–4246. <http://jb.asm.org/content/172/8/4238.long>
- Zhao, J.D. 1989. The Ganodermataceae in China. *Bibliotheca Mycologica* 132: 1–176.

Author contributions: ACB collected the sample, wrote the text and made the analysis; VLB, AMG wrote the text.

Received: 20 January 2016

Accepted: 29 June 2016

Academic editor: Gerardo Robledo